



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/15, A61K 37/64	A1	(11) International Publication Number: WO 91/06650 (43) International Publication Date: 16 May 1991 (16.05.91)
(21) International Application Number: PCT/US90/06072 (22) International Filing Date: 22 October 1990 (22.10.90) (30) Priority data: 428,202 27 October 1989 (27.10.89) US (71) Applicant: CETUS CORPORATION [US/US]; 1400 Fifty-Third Street, Emeryville, CA 94608 (US). (72) Inventors: ELDERING, Eric ; AARDEN, Lucien ; P.O. Box 9190, NL-Amsterdam (NL). (74) Agent: GIOTTA, Gregory, J.; Cetus Corporation, 1400 Fifty-Third Street, Emeryville, CA 94608 (US).		(81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i>
(54) Title: C1 INHIBITOR MUTEINS AND USES THEREOF (57) Abstract Compositions consisting of C1 inhibitor muteins having biological activity similar to C1 inhibitor, but with enhanced resistance to proteolytic cleavage thus rendering such muteins suitable as anti-inflammatory agents, preferably for the treatment or prevention of sepsis.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	SD	Sudan
CF	Central African Republic	KP	Democratic People's Republic of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SN	Senegal
CH	Switzerland	LI	Liechtenstein	SU	Soviet Union
CI	Côte d'Ivoire	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

C1 INHIBITOR MUTEINS AND USES THEREOF

This invention is in the area of molecular biology/immunology, and presents genetically engineered constructs of C1 inhibitor, termed C1 inhibitor muteins, that are resistant to proteolytic attack. The muteins have considerable applications, preferably
5 as anti-inflammatory agents and more preferably for the prophylactic or therapeutic treatment of sepsis.

In the United States alone nosocomial bacteremia develops in about 194,000 patients, and of these about 75,000 die. Maki, D.G., 1981, Nosocomial Infect., (Dikson, R.E., Ed.), page 183, Yrke Medical Books, U.S.A.. Most of these deaths
10 are attributable to six major gram-negative bacilli, and these are *Pseudomonas aeruginosa*, *Escherichia coli*, *Proteus*, *Klebsiella*, *Enterobacter* and *Serratia*. The current treatment for bacteremia is the administration of antibiotics which, unfortunately, have limited effectiveness.

The precise pathology of bacteremia is not completely elucidated, nevertheless,
15 it is known that bacterial endotoxins, lipopolysaccharides (LPS), are the primary causative agent. LPS consist of at least three significant antigenic regions, the lipid A, core polysaccharide, and O-specific polysaccharide. The latter is also referred to as O-specific chain or simply O-antigen. The O-specific chain region is a long-chain polysaccharide built up from repeating polysaccharide units. The number of
20 polysaccharide units differs among different bacterial species and may vary from one to as many as six or seven monosaccharide units. While the O-specific chain varies among different gram-negative bacteria, the lipid A and core polysaccharides are similar if not identical.

Since LPS plays a key role in sepsis, a variety of approaches has been pursued
25 to neutralize its activity. Presently, there is considerable work which suggest that antibody to LPS will soon be a valuable clinical adjunct to the standard antibiotic therapy.

LPS initiates a cascade of biochemical events that eventually causes the death of the patient. It is widely believed that the second event, after the introduction of LPS, is
30 the production of tumor necrosis factor (TNF) as a result of LPS stimulation of macrophage cells. Thus, considerable effort has been expended to produce neutralizing antibody to TNF, or other molecules that could inhibit its septic effects. It is likely that antibody to TNF will have valuable clinical applications. Tracey, et al., 1987, Nature, 330:662.

35 Sepsis caused by gram-negative bacteria is thought to involve activation of the complement system and causes a depletion of various complement component. One

component of a complement system, C5a, causes the aggregation of neutrophils and the aggregates are thought to embolize and cause ischemia. Siegel, J., 1981, Ann. Rev. Med., 32:175. It has been proposed that C5a is thus responsible for the observed organ failure phenomena in sepsis.

5 C1 is a plasma glycoprotein with a molecular weight of about 105,000 and is a member of the super family of serine protease inhibitors which include such members as α 1-antitrypsin, α 1-antiplasmin, antithrombin III, and plasminogen activator inhibitor types I and II. One mechanism by which the activator components of the complement system are controlled is by the C1 inhibitor. The C1 inhibitor is known to
10 inhibit activating components of the classical pathway of complement (C1r and C1s) and the intrinsic coagulation system (Factor XIa, Factor XIIa, and Kallikrein). Further, C1 inhibitor has been shown to interact with the fibrinolytic components plasmin and tissue plasminogen activator.

C1 inhibitor is susceptible to proteolytic cleavage by so called non-target
15 proteases, particularly lysosomal serine protease elastase. Browere, M. and Harpel, P., 1982, J. Biol. Chem., 257:9849. This enzyme is released from polymorphonuclear leukocytes and is present in the circulation of septaremic patients. It is thought that the decrease in the concentration of coagulation factors observed in these patients may, in part, be the result of proteolysis by leukocyte elastase of C1
20 inhibitor. It will be appreciated, that a possible prophylactic/therapeutic approach to treating sepsis would be to genetically engineer C1 inhibitors that are resistant to proteolytic cleavage and administer these to patients that are at risk of contracting sepsis, or that are already septic.

The life threatening nature of sepsis mandates the identification and
25 development of additional therapeutics or prophylactics, both antibody based or otherwise, that may be efficaciously applied in the treatment of sepsis.

In its most general form, the invention described herein presents C1 inhibitor muteins, methods of constructing the muteins, and applications of the muteins, preferably as anti-inflammatory agents and more preferably for the prophylactic or
30 therapeutic treatment of sepsis.

A second object of the invention described herein relates to C1 inhibitor muteins that are both elastase resistant, and that maintain the capacity to covalently bind to, and inactivate components of the complement system.

A third object of the invention is a description of C1 inhibitor muteins that have
35 amino acids at positions 440 and/or 442 mutated to another suitable amino acid, or

deleted, that are both elastase resistant, and that maintain the capacity to covalently bind to, and inactivate components of the complement system.

A fourth object of the invention is a description of C1 inhibitor muteins that display differential sensitivity to proteases and inhibitory activity depending on the type of amino acid that is substituted for amino acids at positions 440 and/or 442.

A fifth object of the invention is a description of C1 inhibitor muteins that exhibit differential inhibitory activity towards various substrates depending on the type of amino acid that is substituted for amino acids at positions 440 and/or 442.

Further, the invention concerns the prophylactic or therapeutic use of C1 inhibitor muteins as anti-inflammatory medicaments, and preferably for the prophylactic or therapeutic treatment of sepsis.

These and further objects of the invention will become apparent after a consideration of the detailed description of the invention shown below.

Figure 1 shows the cDNA sequence corresponding to recombinant C1 inhibitor.

Figure 2 schematically presents a generalized assay procedure for determining the inhibitory or protease sensitivity of the C1 muteins.

Figure 3 shows the degree of inhibitory activity (complex formation) and protease sensitivity (inactivation) of C1 muteins to C1s and Kallikrein.

Figure 4 shows the degree of inhibitory activity (complex formation) and protease sensitivity (inactivation) of C1 muteins to B-12a and plasmin.

Figure 5 shows the amount of neutrophil elastase needed for 50% inhibition of several C1s inhibitor muteins.

1. Definitions

To facilitate understanding the nature and scope of applicant's invention, several definitions regarding various aspects of the invention are presented below. It will be understood, however, that these definitions are general in nature, and encompassed within the definitions are meanings well known to those skilled in the art.

Sepsis is herein defined to mean a disease resulting from gram-positive or gram-negative bacterial infection, the latter primarily due to the bacterial endotoxin, lipopolysaccharide (LPS). It can be induced by at least the six major gram-negative bacilli and these are *Pseudomonas aeruginosa*, *Escherichia coli*, *Proteus*, *Klebsiella*, *Enterobacter* and *Serratia*.

By C1 inhibitor is meant a plasma glycoprotein with a molecular weight of about 105,000 that belongs to the super family of serine protease inhibitors. It inhibits activated components of the classical pathway of complement, C1r and C1s, and the intrinsic coagulation system, factor XIa, factor XIIa, and Kallikrein. C1 also interacts

with plasmin and tissue plasminogen activator. C1 has the further property of itself being inactivated by proteases, notably elastase. It will, of course, be understood that intended to come within the scope of the definition of the C1 inhibitor are fragments of the molecule that maintain biological activity.

5 By C1 inhibitor mutein is meant a molecule that has the biological activity of C1 inhibitor, although to different degrees as exemplified by the data of the invention, and is particularly resistant to proteolytic attack.

Several patents/patent applications and scientific references are referred to below. The instant invention draws on some of the material and methods shown in
10 these references, and thus it is intended that all of the references, in their entirety, be incorporated by reference.

2. C1 Inhibitor Muteins

C1 inhibitor has been cloned and expressed and thus is readily available to the practitioner to perform the herein described mutagenesis. For example, cloning and
15 expression is described by Bock, *et al.*, 1986, Biochemistry, 25:4292. The cDNA sequence is shown in Figure 1. Further, Eldering, *et al.*, 1988, J. Biol. Chem., 263:11776, show a Aat II-HaeII C1 inhibitor cDNA fragment that encodes the entire molecule. This fragment can be manipulated using the procedures described below to generate the C1 inhibitor muteins.

20 A. Mutein Construction--General Procedures

Construction of suitable vectors containing the desired coding and control sequences for the Aat II-HaeII C1 inhibitor cDNA fragment employs standard ligation and restriction techniques which are understood in the art. Isolated plasmids, DNA sequences, or synthesized oligonucleotides are cleaved, tailored, and religated in the form
25 desired.

More specifically, site specific DNA cleavage is performed by treating with the suitable restriction enzyme (or enzymes) under conditions which are generally understood in the art, and the particulars of which are specified by the manufacturer of these commercially available restriction enzymes. See, e.g., New England Biolabs,
30 Product Catalog. In general, about 1 μ g of plasmid or DNA sequence is cleaved by one unit of enzyme in about 20 λ of buffer solution; in the examples herein, typically, an excess of restriction enzyme is used to insure complete digestion of the DNA substrate. Incubation times of about one hour to two hours at about 37°C are workable, although variations can be tolerated. After each incubation, protein is removed by

extraction with phenol/chloroform, and may be followed by ether extraction, and the nucleic acid recovered from aqueous fractions by precipitation with ethanol and resuspension in 10 mM Tris, 1 mM EDTA, pH 7.5. If desired, size separation of the cleaved fragments may be performed by polyacrylamide gel or agarose gel electrophoresis using standard techniques. A general description of size separations is found in Methods in Enzymology, 1980, 65:499-560.

Restriction cleaved fragments may be blunt ended by treating with the large fragment of *E. coli* DNA polymerase I (Klenow) in the presence of the four deoxynucleotide triphosphates (dNTPs) using incubation times of about 15 to 25 minutes at 20 to 25°C in 50 mM Tris pH 7.6, 50 mM NaCl, 6 mM MgCl₂, 6 mM DTT and 5-10 μM dNTPs. The Klenow fragment fills in at 5' sticky ends but chews back protruding 3' single strands, even though the four dNTPs are present. If desired, selective repair can be performed by supplying only one of the, or selected, dNTPs within the limitations dictated by the nature of the sticky ends. After treatment with Klenow, the mixture is extracted with phenol/chloroform and ethanol precipitated followed by running over a Sephadex G-50 spin column. Treatment under appropriate conditions with S1 nuclease results in hydrolysis of any single-stranded portion.

Synthetic oligonucleotides are prepared by the triester method of Matteucci *et al.*, 1981, J. Am. Chem. Soc., 103:3185, or using commercially available automated oligonucleotide synthesizers. Kinasing of single strands prior to annealing or for labelling is achieved using an excess, e.g., approximately 10 units of polynucleotide kinase to 0.1 nmole substrate in the presence of 50 mM Tris, pH 7.6, 10 mM MgCl₂, 5 mM dithiothreitol, 1-2 mM ATP, 1.7 pmoles γ³²P-ATP (2.9 mCi/mmole), 0.1 mM spermidine, 0.1 mM EDTA.

Ligations are performed in 15-30 λ volumes under the following standard conditions and temperatures: 20 mM Tris-Cl pH 7.5, 10 mM MgCl₂, 10 mM DTT, 33 μg/ml BSA, 10 mM-50 mM NaCl, and either 40 μM ATP, 0.01-0.02 (Weiss) units T4 DNA ligase at 0°C (for "sticky end" ligation) or 1 mM ATP, 0.3-0.6 (Weiss) units T4 DNA ligase at 14°C (for "blunt end" ligation). Intermolecular "sticky end" ligations are usually performed at 33-100 μg/ml total DNA concentrations (5-100 nM total end concentration). Intermolecular blunt end ligations (usually employing a 10-30 fold molar excess of linkers) are performed at 1 μM total ends concentration.

In vector construction employing "vector fragments", the vector fragment is commonly treated with bacterial alkaline phosphatase (BAP) in order to remove the 5' phosphate and prevent religation of the vector. BAP digestions are conducted at pH 8 in approximately 150 mM Tris, in the presence of Na⁺ and Mg²⁺ using about 1 unit of

BAP per μg of vector at 60°C for about 1 hour. In order to recover the nucleic acid fragments, the preparation is extracted with phenol/chloroform and ethanol precipitated and desalted by application to a Sephadex G-50 spin column. Alternatively, religation can be prevented in vectors which have been double digested by additional restriction enzyme digestion of the unwanted fragments.

For portions of vectors derived from cDNA or genomic DNA which require sequence modifications, site specific primer directed mutagenesis is used. This is conducted using a synthetic primer oligonucleotide complementary to a single stranded phage DNA to be mutagenized except for limited mismatching, representing the desired mutation. Briefly, the synthetic oligonucleotide is used as a primer to direct synthesis of a strand complementary to the phage, and the resulting double-stranded DNA is transformed into a phage-supporting host bacterium. Cultures of the transformed bacteria are plated in top agar, permitting plaque formation from single cells which harbor the phage.

Theoretically, 50% of the new plaques will contain the phage having, as a single strand, the mutated form; 50% will have the original sequence. The resulting plaques are hybridized with kinased synthetic primer at a temperature which permits hybridization of an exact match, but at which the mismatches with the original strand are sufficient to prevent hybridization. Plaques which hybridize with the probe are then picked, cultured, and the DNA recovered. Details of site specific mutation procedures are described below in specific examples.

Correct ligations for plasmid construction are confirmed by first transforming E. coli strain MM294 obtained from E. coli Genetic Stock Center, CGSC #6135, or other suitable host with the ligation mixture. Successful transformants are selected by ampicillin, tetracycline or other antibiotic resistance or using other markers depending on the mode of plasmid construction, as is understood in the art. Plasmids from the transformants are then prepared according to the method of Clewell, D.B., et al., 1969, Proc. Natl. Acad. Sci. (USA), 62:1159, optionally following chloramphenicol amplification (Clewell, D.B., 1972, J. Bacteriol., 110:667). The isolated DNA is analyzed by restriction and/or sequenced by the dideoxy method of Sanger, F., et al., 1977, Proc. Natl. Acad. Sci. (USA), 74:5463 as further described by Messing et al., 1981, Nucleic Acids Res., 9:309, or by the method of Maxam et al., 1980, Methods in Enzymology, 65:499.

Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described by Cohen, S.N., Proc. Natl. Acad. Sci. (USA) (1972) 69:2110, or the RbCl_2 method described in Maniatis et al., Molecular Cloning: A Laboratory

Manual (1982) Cold Spring Harbor Press, p. 254 was used for procaryotes or other cells which contain substantial cell wall barriers. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and Van der Eb, Virology, 1978, 52:546 is preferred.

5 Host strains used in cloning and expression herein are as follows. For cloning and sequencing, E. coli strain HB101 may be used as the host. For M13 phage recombinants, E. coli strains susceptible to phage infection, such as E. coli K12 strain DG98 are employed. The DG98 strain has been deposited with ATCC 13 July 1984 and has accession number 1965. A preferred expression system for the C1 inhibitor
10 muteins is the COS-1cell /pSVL vector system shown by Eldering et al., 1988, Journal of Biological Chemistry, 263:11776. pSVL (Pharmacia, Uppsala, Sweden) consists of the SV40 origin of replication, the SV40 late promoter, and the VP1 intron in front of a polylinker followed by the SV40 late polyadenylation signal, fused to a pBR322 fragment containing the origin of replication and ampicillin resistance gene.

15 Mutagenesis can be carried out using any number of procedures known in the art. These techniques are described by Smith, 1985, Annual Review of Genetics, 19:423, and modifications of some of the techniques are described in Methods in Enzymology, 154, part E, (eds.) Wu and Grossman (1987), chapters 17, 18, 19, and 20. The preferred procedure is a modification of the Gapped Duplex site-directed
20 mutagenesis method which is described by Kramer, et al., in chapter 17 of volume 154 of Methods in Enzymology, above; and by Kramer et al., 1984, Nucleic Acids Research, 12:9441.

B. Mutein Construction - Preferred Procedures

Conventional M13 mutagenesis methods involve annealing a short synthetic
25 oligonucleotide to single stranded M13 DNA having a cloned target coding sequence that is sought to be mutagenized. The oligonucleotide is almost, but not entirely complementary to the target sequence and has at least one mispaired nucleotide. After the annealing reaction, the remaining portion of the single stranded DNA must be filled in to give heteroduplex DNA that can be transfected into a suitable host cell which
30 allows for the expression of the mutation. In the gapped duplex method, as described by Kramer, et al., in chapter 17 of the Methods in Enzymology, a partial DNA duplex is constructed that has only the target region exposed, unlike the conventional methods which have the target region and the rest of the single stranded M13 DNA exposed. Like the conventional methods, a short oligonucleotide is annealed to the target region,
35 and extended and ligated to produce a heteroduplex. However, because only a small portion of single-stranded DNA is available for hybridization in the gapped duplex

method, the oligonucleotide does not anneal to undesired sites within the M13 genome. This method has the additional advantage of introducing fewer errors during the formation of the heteroduplex since only a very small region of DNA on either side of the target region has to be filled in.

5 More specifically, the gapped duplex method involves cloning the target Aat II-HaeII C1 inhibitor cDNA fragment into an appropriate M13 phage that carries selectable markers, such as, for example, the stop codon amber mutation. The latter allows for negative selection in a host cell that cannot suppress the effects of the mutation. Preferably the phage is M13mp9 which contains two amber codons in
10 critical phage genes. Thus, the sequence that encodes C1 is cloned into M13mp9 amber+, and single stranded DNA is prepared therefrom using standard techniques. Next, double stranded replicative form DNA from M13 GAP, a genetically engineered M13 derivative that lacks the amber codons is cleaved with the appropriate restriction enzyme. The base sequence of M13 GAP is similar to M13mp18, which lacks both the
15 amber codons and the sequence between base pairs 6172 and 6323. This deletion flanks the multiple cloning sites of the M13mp series and generates a unique restriction site. Gapped duplex DNA is formed, using standard DNA/DNA hybridization techniques, consisting of single stranded DNA having the amber codons, and a second strand of DNA from digested M13 GAP lacking both the amber codons and the C1
20 coding sequences. Thus, the only portion of the gapped duplex that is exposed is the C1 target sequence. The desired oligonucleotide(s) is annealed to the gapped duplex DNA, and any remaining gaps filled in with DNA polymerase and the nicks sealed with DNA ligase to produce a heteroduplex. As applied to the instant invention mutagenesis was performed with a mixture of oligonucleotides that code for 16
25 different muteins. The sequence of the degenerate oligonucleotide used to isolate P5 and P3 double mutants and the P5 leucine and valine single muteins is:

5' GCG GGC C(AG) (GC) AGA G (AG) (GC) GGC GGA G 3'

The oligonucleotide is complementary to nucleotides 1412-1433 of Bock *et al.*, above.

30 In addition to the above, several other single P3 muteins were obtained using the following oligonucleotides:

P3-ala 5' GGTGOGGGGOGGCAGAGATGG 3'

P3-gly 5' GGTGOGGGGCTCCAGAGATGG 3'

P3-arg 5' GGGTGOGGGGCTCTAGAGATGGCG 3'

P3-leu 5' GOGGGCCAGAGAGATGG 3'

35 P3-thr 5' GGGTGOGGGGOGGTAGAGATGGCG 3'

The heteroduplex is transfected, preferably into a mismatch repair deficient host, and mixed phage produced. From the mixed phage population, phage carrying

unmutated C1 DNA, which also have the amber mutations, can be selected against by infecting the mixed phage population into a host cell that cannot suppress the amber mutation. Clones can then be screened for phage that carry a C1 mutation, and the molecules sequenced to determine the position of the mutation. Clones were screened
5 using a mixture of kinased degenerate oligonucleotides. Using the foregoing method, 11 C1 inhibitor muteins were produced.

C1 inhibitor mutein DNA fragments were excised from M13 with the appropriate restriction enzymes and cloned into a vector suitable for expression in COS cells. The vector is pSVL and is shown in Bock *et al.*, above. Alternatively pC1-INH,
10 also described by Bock *et al.*, could be used.

SV40-transformed COS-1 monkey cells were grown in Iscove's Modified Dulbecco Cell Culture Medium containing penicillin and streptomycin. The media was supplemented with 10% v/v) heat-inactivated fetal calf serum. Transfection of the cells was performed essentially as described by Luthmann, and Magnusson, 1983, Nucleic
15 Acids Research, 5:1295. This consisted of incubating subconfluent COS-1 cells in the cell culture media for 90 minutes with supercoiled vector encoding C1 inhibitor mutein DNA (5-7.5 µg/ml) and DEAE-dextran (200 µg/ml). After a 90 minute incubation period, the cells were washed twice with cell culture media, and incubated for an additional 2 hours with cell culture media containing 80 µg/ml chloroquine. Two
20 further washes were performed, and the cells incubated with cell culture media supplemented with 10% fetal calf serum. This media was replaced 24 hours later with serum free media. The latter media was harvested after 72 hours, centrifuged to remove cells and debris, and assayed for C1 inhibitor activity, as described below.

C. Assay for C1 Inhibitor Mutein Activities

Figure 2 presents a generalized assay screening format for determining both the inhibitory (complex formation) or protease sensitivity (inactivation) of the C1 muteins. The latter measures inactivation of the C1 inhibitor muteins by non-target proteases. This procedure is also described by Eldering *et al.*, 1988, in J. of Biological Chem.,
25 263:11776. Modifications of these methods are also known in the art.

Generally, inhibitor activity of C1 muteins may be determined by measuring complex formation between the C1 inhibitor and a substrate. The preferred substrates are, of course, C1s, Kallikrein, B-12a, and plasmin. The C1 inhibitor, or C1 inhibitor muteins, inhibit the protease activity of the substrate by forming a covalent bond with the substrate. These molecules were purified by techniques known in the art, or as
30 described by Liebermann, H., *et al.*, 1984, J. Mol. Biol., 177:531 and Nuijens, J., *et al.*, 1987, Immunology, 61:387.

Many procedures are available for measuring the complex resulting from the interaction of the C1 inhibitor mutein and its target proteinase substrate, and include standard immunochemical, radiochemical, or elisa assays. Levin, M., *et al.*, 1983, *J. Biol. Chem.*, 258:6415, Nuijens, J., *et al.*, 1987, *Thromb. Haemost.*, 58:778, de Agostini, 1985, *PNAS*, 82:5190. The procedures generally consist of contacting a C1 inhibitor mutein with a target substrate in solution to permit complex formation to occur, then separating the complex from uncomplexed reactants, and detecting the amount of complex formed. Other assays may be employed whereby the amount of reactants, that is, free C1 inhibitor mutein or target molecule, remaining in the reaction solution are measured after complex formation has occurred.

The preferred assay is a radioimmune assay based on the observation that functional C1 inhibitor muteins bind to activated target substrate, such as C1s. The assay can be performed in several ways, but preferably purified activated C1s is coupled to a solid matrix, preferably Sepharose 4B via cyanogen bromide as is known in the art. Coupled C1s is incubated in an appropriately buffered solution containing, if desired, a small amount of detergent, preferably Tween-20. The latter is used at a concentration of about 0.1% (w/v). The capacity of the C1 inhibitor muteins to complex with the target substrate can be detected using anti-C1 inhibitor mutein antibody. Alternatively, a radiolabelled second antibody can be used to detect the bound first antibody. The antibody may be polyclonal or monoclonal and is incubated for a sufficient time to permit detectible binding of the antibody to the C1 inhibitor to occur. After the appropriate washing steps are conducted whereby non-specifically bound radiolabelled antibody is separated from antibody bound to C1 inhibitor muteins, the amount of radioactivity associated with the latter is determined. In this way, and incorporating the appropriate controls in the assay scheme, the inhibitory capacity of the C1 muteins is determined.

Using similar approaches, the non-target protease sensitivity of the C1 inhibitor muteins may be determined. Numerous assays may be employed that measure the amount of intact C1 inhibitor mutein, or fragments that are derived from the muteins, remaining after exposure to protease. Many procedures are available for measuring the protease sensitivity of the C1 inhibitor muteins, and include standard immunochemical, radiochemical, or elisa assays. A solid phase assay is preferred whereby C1 inhibitor muteins are reacted with non-target protease bound to a solid matrix and the amount of proteolysis of the muteins measured by determining the amount of intact mutein remaining, or preferably by detecting fragments of the mutein. Alternatively, the C1 inhibitor mutein may be bound to a solid support matrix and this material subjected to

proteolysis provided attachment of the mutein does not sterically interfere with protease accessibility to the mutein.

An exemplary non-target protease that cleaves C1 is neutrophil elastase. Thus, this enzyme may be coupled to CNBr treated Sepharose 4B, and incubated with a C1 inhibitor mutein, and the presence of proteolytically cleaved C1 mutein monitored using
5 a number of techniques. The preferred procedure is to pellet the Sepharose-coupled elastase, and measure the presence of cleaved C1 inhibitor in the supernatant using an antibody that recognizes this molecule. Such antibodies are available and are described by Nuijens, *et al.*, 1988, *Blood*, 22:1841. They may be attached to a solid matrix to
10 facilitate separating the cleaved C1 inhibitor mutein from the other reactants. The Sepharose beads containing antibody having bound cleaved C1 mutein inhibitor are spun down, washed and separated from the supernatant containing uncleaved C1 inhibitor or cleaved but unbound fragments. Subsequently, the amount of bound
15 cleaved C1 inhibitor can be determined using a second radiolabelled antibody that recognizes the cleaved molecule. Finally, the amount of radioactivity adherent to the Sepharose beads may be determined as an indication of the amount of cleavage resulting from elastase.

The general procedures for determining both the inhibitory or non-target protease sensitivity of the C1 inhibitor muteins are shown in Figure 1. The procedures
20 involving C1s will be described briefly, the procedures for the other substrates is similar with modifications as noted by Eldering, 1988, *J. Biological Chem.*, 263:11776, and as shown by Nuijens *et al.*, above.

As mentioned above, the radioimmune assay to determine the C1 inhibitor activity of the various muteins is based on the observation that functional C1 inhibitor
25 mutein will bind to activated C1s. Thus, purified activated C1s is coupled to CNBr Sepharose 4B and suspended in phosphate-buffered saline, pH 7.4 10 mM EDTA, and 0.1% (w/v) Tween-20. Next, 0.3 ml of this mixture containing 1.5 µg of activated C1s is incubated with various dilutions of the C1 inhibitor muteins for 5 hours. The
30 Sepharose 4B beads are collected, extensively washed, and complexed C1 inhibitor mutein incubated (>4 hours) with ¹²⁵I-polyclonal anti-C1 antibody. The antibody is described by Hack, *et al.*, above. The Sepharose beads are washed, and bound radioactivity determined using a LKB 1260 multigamma II gamma counter.

Inactivation of the C1 inhibitor muteins is determined as follows. Porcine pancreatic elastase is coupled to Sepharose 4B such that about 3.75 mg of elastase is
35 coupled to 300 mg of Sepharose. The beads are suspended in phosphate buffered saline containing 10 mM EDTA, and 0.1% w/v Tween-20. Various amounts of C1 inhibitor muteins are incubated with 150 µl of Sepharose 4B suspension containing 5.6

mg of elastase in 500 μ l of volume for 1 hour at room temperature. Subsequently, the mixture is centrifuged, and the supernatant assayed for the presence of inactivated C1 inhibitor mutein using KOK12 monoclonal antibody bound to Sepharose, and polyclonal 125 I-anti-C1-inhibitor antibodies, as described above.

5 Figure 3 shows the degree of inhibitory activity (complex formation) and protease sensitivity (inactivation) of 11 C1 inhibitor muteins as assessed using C1s and Kallikrein as substrates and neutrophil elastase as the source of protease. Figure 4 shows the same data with the exception that the substrates were B-12a and plasmin.

10 It is noteworthy that the 11 C1 inhibitor muteins exhibit considerable variation in inhibitor activity, and protease sensitive. This was a function of both the type of amino acid used to substitute for the wild type amino acid, and the substrates used to test for inhibitory activity.

15 The wild type C1 inhibitor has isoleucine and valine at positions 440 and 442, respectively. Mutations at position 440 and 442 are termed P₅ and P₃ muteins, respectively. The figures show that muteins altered only at P₃ (Ala, Gly, Arg, Leu or Thr substituted for Val) display different properties depending on the target protease used. When solid phase C1s is used, complex formation occurs in the order Arg = Gly<Ala<Leu<wild-type = Thr.

20 Further, it appears that two variants, P₅-leu:P₃-ala; and P₅-leu:P₃-leu show significantly reduced susceptibility to inactivation. The amount of HNE needed for 50% inactivation, as determined by residual functional activity towards C1s, is increased by a factor 5 to 8 compared to wild type C1 inhibitor (Figure 5).

D. C1 Pegylated Muteins

25 The preferred embodiment C1 muteins consists of modified muteins that have a substantially longer in vivo circulating lifetime than the unmodified molecules. Favored modified C1 inhibitor muteins are those having a water soluble polymer bound to the muteins. Exemplary of such water soluble polymers are polyacrylic acid, and derivatives thereof, dextran, carboxymethylcellulose, polyethylene glycol, and polyoxyethylated glycerol. The preferred embodiment water soluble polymer is
30 polyethylene glycol. It is disclosed in U.S. Patent No. 4,179,337, along with methods for binding polyethylene glycol to proteins. Polyethylene glycol modified IL-2 is shown in U.S. Patent No. 4,766,106. Using the compositions and procedures described in these two patents, polyethylene glycol modified C1 muteins are readily produced by those skilled in the art.

E. Administration of C1 Muteins

It will be appreciated by those skilled in the art that the C1 muteins described herein can be administered to mammals, including humans, either alone or in combination with other anti-inflammatory agents, or they may be combined with various pharmaceutically acceptable diluents or carriers. Such are widely known to those skilled in the art and are formulated according to standard pharmaceutical practices.

Exemplary diluents include physiologic saline, or buffered saline, as well as Ringer's and dextrose injection fluid, and dextrose saline and lactated Ringer's injection or diluent solutions containing additional therapeutic agents, preferably antibiotics or antibody known to be efficacious in the treatment of sepsis. Such antibody would include those known to be beneficial for the therapeutic treatment of sepsis caused by different strains of bacteria, preferably *Pseudomonas aeruginosa*, *Escherichia coli*, *Proteus*, *Klebsiella*, *Enterobacter* and *Serratia*.

F. Therapeutic Application of C1 Inhibitor Muteins

One embodiment of the invention is the administration of an effective amount of the subject C1 inhibitor muteins to individuals that are at a high risk of developing sepsis or that have developed sepsis. An example of the former category are patients about to undergo surgery. While the mode of administration is not particularly important, parenteral administration is preferred because of the rapid progression of sepsis, and thus, the need to have the C1 inhibitor muteins compositions disseminate quickly throughout the body. The preferred mode of administration is to deliver an I.V. bolus slightly before, during, or after surgery. The dosage of the C1 inhibitors will normally be determined by the prescribing physician. It is to be expected that the dosage will vary according to the age, weight and response of the individual patient.

Having generally described what the applicants believe their invention to be, presented below are examples that are illustrative of the scope of the invention. It will be appreciated by those skilled in the art that the examples are not intended to be construed as limiting the invention to the materials and methods shown as there are numerous substitutions that can be made therein without departing from the scope of the invention.

The present invention has been described with reference to specific embodiments. However, this application is intended to cover those changes and substitutions which may be made by those skilled in the art without departing from the spirit and the scope of the appended claims.

WE CLAIM:

1. Recombinant C1 inhibitor muteins.
2. The C1 inhibitor muteins of claim 1, wherein said muteins are of human origin.
3. The C1 inhibitor muteins of claim 2, wherein said C1 inhibitor muteins have amino acid at position 440 of recombinant C1 inhibitor replaced or deleted.
4. The C1 inhibitor muteins of claim 3, wherein said C1 inhibitor muteins have amino acid at position 440 of recombinant C1 inhibitor replaced with neutral amino acids.
5. The C1 inhibitor muteins of claim 3, wherein said C1 inhibitor muteins have amino acid at position 440 of recombinant C1 inhibitor replaced with charged amino acids.
6. The C1 inhibitor muteins of claim 3, wherein said C1 inhibitor muteins have amino acid at position 440 of recombinant C1 inhibitor replaced with the charged amino acid arginine.
7. The C1 inhibitor muteins of claim 4, wherein said C1 inhibitor muteins have amino acid at position 440 of recombinant C1 inhibitor replaced with neutral amino acids selected from the group consisting of alanine, glycine, leucine, and threonine.
8. The C1 inhibitor muteins of claim 2, wherein said C1 inhibitor muteins have amino acid at position 442 of recombinant C1 inhibitor replaced.
9. The C1 inhibitor muteins of claim 3, wherein said C1 inhibitor muteins have amino acid at position 442 of recombinant C1 inhibitor replaced with neutral amino acids.
10. The C1 inhibitor muteins of claim 3, wherein said C1 inhibitor muteins have amino acid at position 442 of recombinant C1 inhibitor replaced with charged amino acids.

11. The C1 inhibitor muteins of claim 3, wherein said C1 inhibitor muteins have amino acid at position 442 of recombinant C1 inhibitor replaced with the charged amino acid arginine.
12. The C1 inhibitor muteins of claim 4, wherein said C1 inhibitor muteins have amino acid at position 442 of recombinant C1 inhibitor replaced with neutral amino acids selected from the group consisting of alanine, glycine, leucine, and threonine.
13. The C1 inhibitor muteins of claim 2, wherein amino acids at positions 440 and 442 of recombinant C1 inhibitor are replaced by neutral amino acids.
14. The C1 inhibitor muteins of claim 13, wherein amino acids at positions 440 and 442 of recombinant C1 inhibitor are replaced by neutral amino acid alanine and leucine, respectively.
15. The C1 inhibitor muteins of claim 13, wherein amino acids at positions 440 and 442 of recombinant C1 inhibitor are replaced by neutral amino acids alanine and valine, respectively.
16. The C1 inhibitor muteins of claim 13, wherein amino acids at positions 440 and 442 of recombinant C1 inhibitor are replaced by neutral amino acid leucine.
17. The C1 inhibitor muteins of claim 13, wherein amino acids at positions 440 and 442 of recombinant C1 inhibitor are replaced by neutral amino acids leucine and valine, respectively.
18. Recombinant DNA that encodes a molecule comprising C1 inhibitor mutein activity.
19. Recombinant DNA that encodes a molecule comprising C1 inhibitor mutein activity as described in claim 7.
20. Recombinant DNA that encodes a molecule comprising C1 inhibitor mutein activity as described in claim 12.

21. Recombinant DNA that encodes a molecule comprising C1 inhibitor mutein activity as described in claim 17.
22. A composition for the therapeutic or prophylactic treatment of sepsis comprising an effective amount of a biologically active C1 inhibitor mutein as described in claim 7.
23. A composition for the therapeutic or prophylactic treatment of sepsis comprising an effective amount of a biologically active C1 inhibitor mutein as described in claim 12.
24. A composition for the therapeutic or prophylactic treatment of sepsis comprising an effective amount of a biologically active C1 inhibitor mutein as described in claim 17.
25. A method for treating sepsis comprising administering an effective amount of the composition of claim 22 to a patient.
26. A method for treating sepsis comprising administering an effective amount of the composition of claim 23 to a patient.
27. A method for treating sepsis comprising administering an effective amount of the composition of claim 24 to a patient.

1 / 8

CCAGAAGTTTGGAGTCCGCTGACGTCGCCGCCAG ATG GCC TCC AGG CTG ACC
 -22 -20
 met ala ser arg leu thr

TCA AAT CCA AAT GCT ACC AGC TCC AGC TCC CAG GAT CCA GAG AGT
 1 10
 ser asn pro asn ala thr ser ser ser ser gln asp pro glu ser

I
 Y
 S S

AAG ATG CTA TTC GTT GAA CCC ATC CTG GAG GTT TCC AGC TTG CCG
 30 40
 lys met leu phe val glu pro ile leu glu val ser ser leu pro

CE
 T M

ACC ACT GAT GAA CCC ACC ACA CAA CCC ACC ACA GAG CCC ACC ACC
 60 70
 thr thr asp glu pro thr thr gln pro thr thr glu pro thr thr

GAT TCT CCT ACC CAG CCC ACT ACT GGG TCC TTC TGC CCA GGA CCT
 90 100
 asp ser pro thr gln pro thr thr gly ser phe cys pro gly pro

CE
 TTG GGG GAT GCT TTG GTA GAT TTC TCC CTG AAG CTC TAC CAC GCC
 120 130
 leu gly asp ala leu val asp phe ser leu lys leu tyr his ala

TTC AGC ATC GCC AGC CTC CTT ACC CAG GTC CTG CTC GGG GCT GGG
 150 160
 phe ser ile ala ser leu leu thr gln val leu leu gly ala gly

TC
 GAC TTC ACC TGT GTC CAC CAG GCC CTG AAG GGC TTC ACG ACC AAA
 180 190
 asp phe thr cys val his gln ala leu lys gly phe thr thr lys

C T C
 ATA AGG GAC ACC TTT GTG AAT GCC TCT CGG ACC CTG TAC AGC AGC

FIG. 1A

2 / 8

CTG CTG ACC CTC CTG CTG CTG CTG CTG GCT GGG GAT AGA GCC TCC 98
 leu leu thr leu leu leu ⁻¹⁰ leu leu leu ala gly asp arg ala ser
 TTG CAA GAC AGA GGC GAA GGG AAG GTC GCA ACA ACA GTT ATC TCC 188
 leu gln asp arg gly ²⁰ glu gly lys val ala thr [◇] thr val ile ser
 ACA ACC AAC TCA ACA ACC AAT TCA GCC ACC AAA ATA ACA GCT AAT 278
 thr thr [◇] asn ser [◇] thr ⁵⁰ thr asn ser ala thr lys ile thr ala [◇] asn
 CAA CCC ACC ATC CAA CCC ACC CAA CCA ACT ACC CAG CTC CCA ACA 368
 gln pro thr ⁸⁰ ile gln pro thr gln pro thr thr gln leu pro thr
 GTT ACT CTC TGC TCT GAC TTG GAG AGT CAT TCA ACA GAG GCC GTG 458
 val thr leu ¹¹⁰ (cys) ser asp leu glu ser his ser thr glu ala val
 TTC TCA GCA ATG AAG AAG GTG GAG ACC AAC ATG GCC TTT TCC CCA 548
 phe ser ala met lys ¹⁴⁰ lys val glu thr asn met ala phe ser pro
 CAG AAC ACC AAA ACA AAC CTG GAG AGC ATC CTC TCT TAC CCC AAG 630
 gln asn thr lys thr ¹⁷⁰ asn leu glu ser ile leu ser tyr pro lys
 GGT GTC ACC TCA GTC TCT CAG ATC TTC CAC AGC CCA GAC CTG GCC 728
 gly val thr ser val ²⁰⁰ ser gln ile phe his ser pro asp leu ala
 AGC CCC AGA GTC CTA AGC AAC AAC AGT GAC GCC AAC TTG GAG CTC 818

FIG. 1B

210 ile arg asp thr phe val asn ala ser arg thr leu tyr ser ser
ATC AAC ACC TGG GTG GCC AAG AAC ACC AAC AAC AAG ATC AGC CGG
240 ile asn thr trp val ala lys asn thr asn asn lys ile ser arg
GCT ATC TAC CTG AGT GCC AAG TGG AAG ACA ACA TTT GAT CCC AAG
270 ala ile tyr leu ser ala lys trp lys thr thr phe asp pro lys
GTG CCC ATG ATG AAT AGC AAG AAG TAC CCT GTG GCC CAT TTC ATT
300 val pro met met asn ser lys lys tyr pro val ala his phe ile
AAT CTG AGT TTG GTG ATC CTG GTA CCC CAG AAC CTG AAA CAT CGT
asn leu ser leu val ile leu val pro gln asn leu lys his arg
ATC ATG GAG AAA CTG GAG ATG TCC AAG TTC CAG CCC ACT CTC CTA
360 ile met glu lys leu glu met ser lys phe gln pro thr leu leu
ATC ATG GAG AAA TTG GAA TTC TTC GAT TTT TCT TAT GAC CTT AAC
390 ile met glu lys leu glu phe phe asp phe ser tyr asp leu asn
CAG CAC CAG ACA GTG CTG GAA CTG ACA GAG ACT GGG GTG GAG GCG
420 gln his gln thr val leu glu leu thr glu thr gly val glu ala
GAA GTG CAG CAG CCC TTC CTC TTC GTG CTC TGG GAC CAG CAG CAC
450 glu val gln gln pro phe leu phe val leu trp asp gln gln his
GACCTGCAGGATCAGGTTAGGGCGAGCGCTACCTCTCCAGCCTCAGCTCTCAGTTGCAGC
CCAAAAGGGCTCCTGAGGGTCTGGGCAAGGGACCTGCTTCTATTAGCCCTTCTCCATGGC
ATA AATAAA ACCTGACAGACCATAAAAAAAAAA

FIG. 1C

FIG. 1C




FIG. 1D

5 / 8

Schematic Representation of RIAs

- target protease (C1S etc...) or non-target
- protease (elastase) coupled to Sepharose
- add rC1 inhibitor or muteins (serial dilutions)
- >4 hour incubation
- spin down Sepharose

Pellet - Assay for Complex Formation

Supernatant - Assay for Elastase Sensitivity

- 5 x wash
- >4 hour incubation
- add poly-a-C1 Inh*
- 6 x wash
- count bound radioactivity


- add mAb KOK12 Sepharose
(detects cleaved C1 Inh.)
- >4 hour incubation
- 5 x wash
- add poly-a-C1 Inh*
- >4 hour incubation
- 5 x wash
- count bound radioactivity

FIG. 2

6 / 8


CIS

FIG. 3A




KALLIKREIN

FIG. 3B




7 / 8

B-12a


FIG. 4A**FIG. 4B**

PLASMIN



A DASH (-) MEANS WILD TYPE RESIDUE.

FIG. 5



INTERNATIONAL SEARCH REPORT

International Application No PCT/US 90/06072

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶ According to International Patent Classification (IPC) or to both National Classification and IPC IPC ⁵ : C 12 N 15 /15, A 61 K 37/64										
II. FIELDS SEARCHED <div style="text-align: right; font-size: small;">Minimum Documentation Searched ⁷</div> <table style="width: 100%; border: none;"> <tr> <td style="width: 25%; border-bottom: 1px solid black; font-size: small;">Classification System</td> <td style="border-bottom: 1px solid black; font-size: small;">Classification Symbols</td> </tr> <tr> <td style="height: 40px; vertical-align: top; border: none;">IPC⁵</td> <td style="height: 40px; vertical-align: top; border: none;">C 12 N</td> </tr> </table> <div style="text-align: center; font-size: x-small; margin-top: 10px;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸</div>			Classification System	Classification Symbols	IPC ⁵	C 12 N				
Classification System	Classification Symbols									
IPC ⁵	C 12 N									
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹ <table border="1" style="width: 100%; border-collapse: collapse; font-size: small;"> <tr> <th style="width: 10%;">Category [*]</th> <th style="width: 70%;">Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²</th> <th style="width: 20%;">Relevant to Claim No. ¹³</th> </tr> <tr> <td style="vertical-align: top; padding: 5px;">O, X</td> <td style="vertical-align: top; padding: 5px;"> Complement Inflammation, vol. 6, no. 5, XIIIth International Complement Workshop, 10-15 September 1989, San Diego, California, E. Eldering et al.: "C1 inhibitor variants with reduced susceptibility for neutrophil elastase", see page 333, abstract 68 <div style="text-align: center; border-top: 1px dashed black; width: 100%; margin: 10px 0;"></div> </td> <td style="vertical-align: top; padding: 5px;"> 1-4, 7-9, 12-14, 18-20, 22-23 </td> </tr> </table>			Category [*]	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³	O, X	Complement Inflammation, vol. 6, no. 5, XIIIth International Complement Workshop, 10-15 September 1989, San Diego, California, E. Eldering et al.: "C1 inhibitor variants with reduced susceptibility for neutrophil elastase", see page 333, abstract 68 <div style="text-align: center; border-top: 1px dashed black; width: 100%; margin: 10px 0;"></div>	1-4, 7-9, 12-14, 18-20, 22-23		
Category [*]	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³								
O, X	Complement Inflammation, vol. 6, no. 5, XIIIth International Complement Workshop, 10-15 September 1989, San Diego, California, E. Eldering et al.: "C1 inhibitor variants with reduced susceptibility for neutrophil elastase", see page 333, abstract 68 <div style="text-align: center; border-top: 1px dashed black; width: 100%; margin: 10px 0;"></div>	1-4, 7-9, 12-14, 18-20, 22-23								
<div style="display: flex; justify-content: space-between; font-size: x-small;"> <div style="width: 45%;"> <p>[*] Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p> </div> </div>										
IV. CERTIFICATION <table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border-bottom: 1px solid black; font-size: small;">Date of the Actual Completion of the International Search</td> <td style="width: 50%; border-bottom: 1px solid black; font-size: small;">Date of Mailing of this International Search Report</td> </tr> <tr> <td style="height: 40px; vertical-align: bottom; text-align: center;">23rd January 1991</td> <td style="height: 40px; vertical-align: bottom; text-align: center;">1 MAR 1991</td> </tr> <tr> <td style="border-bottom: 1px solid black; font-size: small;">International Searching Authority</td> <td style="border-bottom: 1px solid black; font-size: small;">Signature of Authorized Officer</td> </tr> <tr> <td style="height: 40px; vertical-align: bottom; text-align: center;">EUROPEAN PATENT OFFICE</td> <td style="height: 40px; vertical-align: bottom; text-align: center;"> MISS D </td> </tr> </table>			Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	23rd January 1991	1 MAR 1991	International Searching Authority	Signature of Authorized Officer	EUROPEAN PATENT OFFICE	 MISS D
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report									
23rd January 1991	1 MAR 1991									
International Searching Authority	Signature of Authorized Officer									
EUROPEAN PATENT OFFICE	 MISS D									

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE ¹

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☒ Claim numbers ~~XX~~ because they relate to subject matter not required to be searched by this Authority, namely:

~~XX~~ Claims 25-27

See PCT Rule 39.1(iv)

Methods for treatment of the human or animal body by means of surgery or therapy, as well as diagnostic methods.

2. ☐ Claim numbers because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claim numbers because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING ²

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.